

Non-Gaussianity in the unified curvaton mechanism : The generalized curvaton mechanism that comprehends modulation at the transition

Seishi Enomoto

Department of Physics, Nagoya University, Nagoya 464-8602, Japan

Kazunori Kohri

*Cosmophysics group, Theory Center, IPNS, KEK,
and The Graduate University for Advanced Study (Sokendai), Tsukuba 305-0801, Japan*

Tomohiro Matsuda

*Department of Physics, Lancaster University, Lancaster LA1 4YB, UK,
and Laboratory of Physics, Saitama Institute of Technology, Fukaya, Saitama 369-0293, Japan*

Generation of the curvature perturbation is calculated when the modulation is implemented in the generalized curvaton mechanism, in which the curvaton may not scale like matter. We first consider the slow-roll curvaton scenario with/without modulation at the end of the slow-roll, where the curvaton and the modulation share the same source of the perturbation. We calculate the non-linearity parameter using the non-linear formalism, which is the first exact analytical calculation of the non-Gaussianity created by the slow-roll curvaton. Unlike the conventional curvaton mechanism, in which f_{NL} can become large but arbitrary, our result shows that $f_{NL} \sim O(10)$ could be natural in the typical inflating curvaton scenario. Our calculation is also valid in the conventional modulation that is usually caused by an extra light field, in which the curvaton and the modulation may have the individual (separable) source of the perturbations. For the separable perturbations we consider the simplest multi-field inflation.

PACS numbers: 98.80Cq

I. INTRODUCTION

The primordial curvature perturbation $\zeta(k)$ that exists on cosmological scales just before they start to enter the horizon is usually related to the perturbations generated during inflation. Recent observation suggests that $\zeta(k)$ is strongly constrained and provides a window on the very early universe [1].

The mechanism of generating ζ can be diverse, but begins presumably during inflation. There are many proposals for generating the curvature perturbation from the field perturbations, which use one or more fields for the mechanism.

The paradigm of the multi-field inflation scenario has been widely investigated, but it has usually been supposed that $\zeta(x, t)$ evaluated at an epoch t_{end} just after the end of inflation is to be identified with the observed quantity. In this respect, a lot of papers consider the calculation of the spectrum of ζ just at the end of inflation [2–9].

On the other hand, we know that multi-field inflation always lead to the multi-component Universe after inflation, in which there could be significant evolution of the curvature perturbation after inflation [10]. The typical example is the curvaton mechanism, in which the mixed state of the radiation and the matter causes significant evolution of the curvature perturbation [11, 12]. Recently, the curvaton mechanism has been extended to include the slow-rolling field [13]. And more recently, it has been shown that the curvaton mechanism works in the conventional multi-field inflation without introduc-

ing an extra light field as the curvaton [10]. These recent works show that the evolution “after” multi-field inflation is more diverse than it has been expected, and that the mechanisms will be practically important for the creation of the perturbations of the Universe.

In this paper we go further and point out that the actual calculation of the curvaton mechanism might well depend on the modulated transition [14]¹, *even if there is no moduli that causes conventional “modulation”*. When it is “modulated”, the hypersurface of the transition does not coincide with the uniform density hypersurfaces. The typical example of this kind is the modulated reheating scenario [15], although the original model considers only the single-component Universe. Considering the modulation in the multi-component Universe, and also including a slow-rolling component, we find that the modulation is sometimes caused by the conventional isocurvature perturbations that have been used for the curvaton mechanism.

Moreover, given that multi-field inflation (e.g, N-flaton [16]) is general, and it usually ends with the multi-component Universe, any kind of modulation must be extended to the multi-component Universe, where the curvaton mechanism is important in the same way. In that case, one needs to consider both curvaton and mod-

¹ Sometimes the curvaton mechanism assumes sudden-decay approximation. In this paper, abrupt change of the density scaling relation is assumed for the “transition”.

ulation at the same time.

We basically use the δN formalism for the calculation. In this formalism, ζ is defined by smoothing the energy density ρ on a super-horizon scale shorter than any scale of interest. Then the local energy continuity equation is given by

$$\frac{\partial \rho(x, t)}{\partial t} = -\frac{3}{a(x, t)} \frac{\partial a(x, t)}{\partial t} (\rho(x, t) + p(x, t)), \quad (1)$$

where t is time along a comoving thread of spacetime and a is the scale factor.

During inflation, the vacuum fluctuation of a light scalar field ϕ_i is converted at horizon exit to a nearly Gaussian classical perturbation with spectrum $H/2\pi$. Here $H \equiv \dot{a}(t)/a(t)$ is the Hubble parameter defined in the unperturbed universe. The δN formalism gives

$$\zeta = \delta[\ln(a(x, t)/a(t_*))] \equiv \delta N, \quad (2)$$

where taking t_* to be an epoch during inflation after relevant scales leave the horizon, one can assume $N(\phi_1(x, t_*), \phi_2(x, t_*), \dots, t, t_*)$ so that

$$\zeta(x, t) = N_i \delta \phi_i(x, t_*) + \frac{1}{2} N_{ij} \delta \phi_j(x, t_*) + \dots, \quad (3)$$

where a subscript i denotes the derivative with respect to ϕ_i , which is evaluated on the unperturbed trajectory.

In the curvaton calculation [11, 17] one usually assumes that these expressions are dominated by the single “curvaton” field σ , which starts oscillating in the radiation dominated Universe when σ has the negligible contribution to the curvature perturbation. Then the non-Gaussianity parameter is given by [18, 19]

$$f_{NL} \simeq \frac{5}{4r_\sigma} \left(1 + \frac{g''g}{g'^2} \right) - \frac{5}{3} - \frac{5}{6} r_\sigma, \quad (4)$$

where $g(\sigma_*)$ is the initial amplitude of the oscillation as a function of the curvaton field at horizon exit [18], and r_σ is defined as r_1 in the later paragraph. The above result is obtained for the two-component Universe, in which one component behaves like matter while the other behaves like radiation. *Our calculation is aimed to generalize and extend the above results including any kind of fluid for the components.*

In this paper we consider the non-linear formulation defined in Ref. [20, 21];

$$\begin{aligned} \zeta_i &= \delta N + \int_{\bar{\rho}_i}^{\rho_i} \frac{d\bar{\rho}_i}{3(1+w_i)\bar{\rho}_i} \\ &= \delta N + \frac{1}{3(1+w_i)} \ln \left(\frac{\rho_i}{\bar{\rho}_i} \right) \\ &\simeq \delta N + \frac{1}{3(1+w_i)} \frac{\delta \rho_i^{\text{iso}}}{\bar{\rho}_i}, \end{aligned} \quad (5)$$

where $\delta \rho_i^{\text{iso}}$ will be defined in Eq.(7). Here $w_i = 1/3$ for the radiation fluid and $w_i = 0$ for the matter fluid. A

bar is for the homogeneous quantity, and ρ_i is defined on the uniform density hypersurfaces. Due to the isocurvature perturbations, $\bar{\rho}_i$ and ρ_i are different in the multi-component Universe. The curvature perturbation of the total fluid should be discriminated from the component curvature perturbation ζ_i . The standard definition of the adiabatic perturbation is given by

$$\delta N = -H \frac{\delta \rho^{\text{adi}}}{\dot{\rho}}, \quad (6)$$

where $\delta \rho^{\text{adi}} \equiv \sum_i \delta \rho_i^{\text{adi}}$ must be evaluated on the spatially flat hypersurfaces. In contrast to $\delta \rho_i^{\text{adi}}$, the isocurvature quantity $\delta \rho_i^{\text{iso}}$ is related to the fraction perturbation defined on the uniform density hypersurfaces. Using the homogeneous density $\bar{\rho}_i$, the isocurvature density perturbation is defined on the uniform density hypersurfaces as

$$\delta \rho_i^{\text{iso}} \equiv \rho_i - \bar{\rho}_i, \quad (7)$$

which satisfies $\sum_i \delta \rho_i^{\text{iso}} = 0$. In the multi-component Universe, the uniform density hypersurface defining uniform $\rho \equiv \sum_i \rho_i$ is usually different from the one defining the uniform ρ_i hypersurfaces for each component.

We find from the second line of Eq.(5);

$$\rho_i = \bar{\rho}_i e^{3(1+w_i)(\zeta_i - \delta N)}. \quad (8)$$

Using the above equation, the definition of the total energy density $\rho^{\text{total}} \equiv \rho_1 + \rho_2 = \bar{\rho}_1 + \bar{\rho}_2$ on the uniform density hypersurfaces leads to

$$f_1 e^{3(1+w_1)(\zeta_1 - \delta N)} + (1 - f_1) e^{3(1+w_2)(\zeta_2 - \delta N)} = 1, \quad (9)$$

where the fraction of the energy density is defined by

$$f_1 \equiv \frac{\bar{\rho}_1}{\bar{\rho}_1 + \bar{\rho}_2}. \quad (10)$$

Expanding Eq.(9) and solving the equation for δN , we find at first order

$$\begin{aligned} \delta N &= r_1 \zeta_1 + (1 - r_1) \zeta_2 \\ &= [r_1 \zeta_1^{\text{iso}} + (1 - r_1) \zeta_2^{\text{iso}}] + \zeta^{\text{adi}}, \end{aligned} \quad (11)$$

where $\zeta_i^{\text{iso}} \equiv \zeta_i - \zeta^{\text{adi}}$ is introduced in the last line. r_1 is defined by

$$r_1 \equiv \frac{3(1+w_1)\bar{\rho}_1}{3(1+w_1)\bar{\rho}_1 + 3(1+w_2)\bar{\rho}_2}. \quad (12)$$

Note that ζ^{adi} is identical to δN at the time when the initial quantities are evaluated. Usual curvaton scenario assumes $\delta N = \zeta^{\text{adi}} = 0$ at the beginning and it defines the initial condition. ζ_i is constant in the usual curvaton scenario [20]; in that case both ζ_i^{iso} and ζ^{adi} are constant in the above definition. δN is time-dependent when $r_1(t)$ is changing, which defines the curvaton mechanism. The parameter r_1 is important for the non-Gaussianity. $r_1 \sim 0.1$ will be justified later in this paper.

The definition given by Eq.(11) can be used anytime and for any long-wavelength perturbations. However, because we are formulating the evolution of the adiabatic perturbation that is caused by the adiabatic-isocurvature mixings, we need first to define the “starting point” at an epoch and then we can discuss the “evolution” thereafter. The quantities evaluated at the starting point is denoted by the subscript “ini”.

When we calculate the evolution of the curvature perturbation using the curvaton mechanism, ζ^{adi} and ζ_i are evaluated at the “starting point” and ζ_i is supposed to be constant thereafter. δN in Eq.(11) is identical to ζ^{adi} at the “starting point”, but apart from that point the evolution causes $\delta N \neq \zeta^{\text{adi}}$. This gives the basic mechanism of the generalized curvaton mechanism, which is valid even if the curvaton is not scaling like matter.

The primordial perturbations are usually evaluated at the end of the primordial inflation ($t = t_{\text{end}}$), while the corresponding scale exits horizon at $t_* < t_{\text{end}}$. Using ρ_i evaluated at the end of inflation, ζ_i is given by

$$\begin{aligned}\zeta_{i,\text{end}} &= \zeta^{\text{inf}} + \frac{1}{3(1+w_{i,\text{end}})} \ln \left(\frac{\rho_{i,\text{end}}}{\bar{\rho}_{i,\text{end}}} \right) \\ &\equiv \zeta^{\text{inf}} + \zeta_{i,\text{end}}^{\text{iso}},\end{aligned}\quad (13)$$

where ζ^{inf} is the primordial curvature perturbation that is created during the primordial inflation. Here ζ^{inf} , $\zeta_{i,\text{end}}$, $\zeta_{i,\text{end}}^{\text{iso}}$ and $\rho_{i,\text{end}}$ are evaluated at $t = t_{\text{end}}$.

Later in this paper we consider the secondary inflation for the curvaton mechanism (the inflating curvaton). In that case the end of the secondary inflation is denoted by the subscript “e”, which should be discriminated from the subscript “end” used above.

In the standard curvaton scenario the transition at the end (i.e, the curvaton decay) usually occurs on the uniform density hypersurfaces. In that case there is no modulation, and the creation of the curvature perturbation is simply due to the curvaton mechanism.

On the other hand, in the standard modulation scenario (the modulated reheating [15] and the modulated end of inflation [22]), the transition is modulated but there is only one component in the Universe.

In this paper we are considering the unification of the two scenarios. In the original modulation scenario the source of the modulation is supposed to be the extra light field (moduli), but the source of the modulation could be diverse when the modulation is implemented in the generalized curvaton mechanism. *In this paper we show that the modulation in the generalized curvaton scenario can be sourced by the conventional isocurvature perturbation that is identical to the source of the curvaton mechanism. The new modulation scenario is discussed for the generalized curvaton model using the unified non-linear formalism.*




FIG. 1: Evolution of the density $\rho_1(\rho_2)$ is plotted with the three phases (A, B, C). Typical modulation scenarios are shown in the figure.

II. THE GENERALIZED CURVATON SCENARIO WITHOUT MODULATION

A. Inflating Curvaton

In this section ϕ_1 is the curvaton and ϕ_2 is the inflaton of the primordial inflation. For simplicity, we assume that ϕ_2 decays instantly just after inflation, while ϕ_1 decays late at $H_{d1} \ll H_I$. There is no mixing between ϕ_1 and ϕ_2 . Here H_I denotes the Hubble parameter during the primordial inflation. The sinusoidal ϕ_1 -oscillation of the curvaton field starts at H_{osc} .

Then, there are three phases (A, B, C) characterized by the parameter w_1 , which are separated by $H = H_{\text{osc}}$ and H_{d1} ;

- (A) ϕ_1 ; slow-roll, ρ_2 ; radiation
($1 + w_1 \equiv \epsilon_w \ll 1$, $w_2 = 1/3$)
- (B) ϕ_1 ; oscillating, ρ_2 ; radiation
($w_1 = 0$, $w_2 = 1/3$)
- (C) ρ_1 ; radiation, ρ_2 ; radiation
($w_1 = 1/3$, $w_2 = 1/3$),

where $\epsilon_w \ll 1$ is not always a constant [13]. In contrast to ζ_1 , which changes time to time, ζ_2 is always invariant during the evolution. (See also Fig.1)

Basically, the evolution of the curvature perturbations during the slow-roll phase (A) is not simple, because ζ_1 is not constant in this phase. We thus need further simplifications of the scenario;

1. As far as ρ_2 is dominating the Universe, we always find $r_1 \sim \epsilon_w \frac{\rho_1}{\rho_2} < \epsilon_w \ll 1$. During this period the curvaton mechanism is negligible.

2. The curvaton mechanism becomes significant when ρ_1 starts dominating the Universe (the inflating curvaton). Note however that r_1 is still small ($r_1 \sim \epsilon_w \ll 1$) at the beginning of the curvaton inflation.
3. ζ_1 is not always constant but behaves like constant somewhat after the beginning of the curvaton inflation [13]. In that case the initial (starting-point) perturbation can be calculated at that moment. At the end of the curvaton mechanism (i.e., after the significant evolution of the curvature perturbation) one will find $r_1 \gg \epsilon_w$.

Just after the beginning of the curvaton inflation, we define the initial perturbations

$$\zeta_{1,\text{ini}} \equiv \zeta_{1,\text{ini}}^{\text{adi}} + \zeta_{1,\text{ini}}^{\text{iso}}. \quad (14)$$

Using the same argument as in Eq.(11), we find the curvature perturbation just before the end of the curvaton inflation;

$$\begin{aligned} \delta N_- &\equiv r_{1-}\zeta_{1-} + (1-r_{1-})\zeta_{2-} \\ &\simeq r_{1-}\zeta_{1,\text{ini}} + (1-r_{1-})\zeta_{2,\text{ini}} \\ &= \zeta_{1,\text{ini}}^{\text{adi}} + [r_{1-}\zeta_{1,\text{ini}}^{\text{iso}} + (1-r_{1-})\zeta_{2,\text{ini}}^{\text{iso}}], \end{aligned} \quad (15)$$

where

$$r_{1-} \equiv \frac{3\epsilon_w \bar{\rho}_{1-}}{3\epsilon_w \bar{\rho}_{1-} + 4\bar{\rho}_{2-}}. \quad (16)$$

Here the minus sign denotes the quantities just before the transition. The quantities satisfy $\bar{\rho}_{1-} \simeq \bar{\rho}_{1,\text{ini}}$ and $\bar{\rho}_{1,\text{ini}} \simeq \bar{\rho}_{2,\text{ini}} \gg \bar{\rho}_{2-}$ for the inflating curvaton.

Then, we find the curvature perturbation created by the evolution [13]².

$$\begin{aligned} \delta N &\simeq r_{1-}\zeta_{1,\text{ini}}^{\text{iso}} \\ &\simeq \frac{\delta \rho_{1,\text{ini}}^{\text{iso}}}{3\epsilon_w \bar{\rho}_{1-} + 4\bar{\rho}_{2-}}. \end{aligned} \quad (17)$$

For the curvaton inflation that starts at $\rho_1 \simeq \rho_2$, we find

$$\begin{aligned} \tilde{N}_e &= \ln \left(\frac{\bar{\rho}_{1-}}{\bar{\rho}_{2-}} \right)^{1/4} \\ &= -\frac{1}{4} \left[\ln \epsilon_w + \ln \left(\frac{1-r_{1-}}{r_{1-}} \right) - \ln \frac{4}{3} \right], \end{aligned} \quad (18)$$

where $\tilde{N}_e$ is the number of e-foldings elapsed during the curvaton inflation and we find from Eq.(16);

$$\bar{\rho}_{1-} = \frac{4r_{1-}}{3\epsilon_w(1-r_{1-})} \bar{\rho}_{2-}. \quad (19)$$

If the curvaton inflation is very short, we find that the denominator of Eq.(17) gives $3\epsilon_w \bar{\rho}_{1-} + 4\bar{\rho}_{2-} \simeq 4\bar{\rho}_{2-}$. In that specific case we find

$$\begin{aligned} \delta N^{\text{short}} &\simeq \frac{1}{4} \frac{\delta \rho_{1,\text{ini}}^{\text{iso}}}{\bar{\rho}_{2-}} \\ &\simeq \frac{r_{1-}}{3\epsilon_w} \frac{\delta \rho_{1,\text{ini}}^{\text{iso}}}{\bar{\rho}_{1,\text{ini}}}, \end{aligned} \quad (20)$$

while after enough e-foldings we find

$$\delta N^{\text{long}} \simeq \frac{1}{3\epsilon_w} \frac{\delta \rho_{1,\text{ini}}^{\text{iso}}}{\bar{\rho}_{1,\text{ini}}}. \quad (21)$$

Obviously, the curvature perturbation created during the evolution satisfies $\delta N^{\text{short}} \ll \delta N^{\text{long}}$, which shows that longer inflation (curvaton inflation) is more efficient for the curvaton mechanism. Note however the above result is valid only when the end of the curvaton slow-roll coincides with the uniform density hypersurfaces; otherwise the modulation at the end may not be negligible.

Later in this paper we will show how the significant modulation is possible when the slow-roll ends without inflation.

B. More on non-Gaussianity in the generalized curvaton mechanism

In this paragraph we are basically following the non-linear calculation in Ref.[19], extending the calculation to include the slow-rolling curvaton. Generally one can expand

$$\phi_1 = \bar{\phi} + \sum_{k=1}^{\infty} \frac{1}{k!} \delta^{(k)} \phi_1, \quad (22)$$

where $\delta^{(1)} \phi_1$ is a Gaussian random field. The primordial perturbation can be expanded as

$$\zeta_1 = \zeta_1^{(1)} + \sum_{k=2}^{\infty} \frac{1}{k!} \zeta_1^{(k)}, \quad (23)$$

where $\zeta_1^{(1)}$ is Gaussian. Non-linearity parameters are defined for the adiabatic perturbation ζ ;

$$\zeta = \zeta^{(1)} + \frac{3}{5} f_{NL} (\zeta^{(1)})^2 + \frac{9}{25} g_{NL} (\zeta^{(1)})^3 + \dots \quad (24)$$

Assume that the curvaton potential during the curvaton inflation is approximated by

$$\rho_1 = V_0 \pm \frac{1}{2} m^2 \phi_1^2, \quad (25)$$

where the effective mass term may have either positive or negative sign.

² The standard assumptions of the curvaton mechanism are $\zeta_2 \ll \zeta_1$ and $\zeta_1^{\text{inf}} \simeq 0$. The above result is obtained using these assumptions.

Using the Gaussian quantum fluctuations at the horizon exit ($\delta\phi_{1*}$), we can write [19]

$$\phi_{1*} = \bar{\phi}_{1*} + \delta\phi_{1*}. \quad (26)$$

In that case we write

$$\phi_{1,\text{ini}} \equiv g(\phi_{1*}) \quad (27)$$

and expand [19]

$$\phi_{1,\text{ini}} = \bar{g} + \sum_{k=1}^{\infty} \frac{1}{k!} g^{(k)} \left(\frac{\bar{g}}{g'} \frac{\delta\phi_1}{\bar{\phi}_1} \right)^k, \quad (28)$$

where we wrote $g^{(n)} \equiv \partial^n g / \partial \phi_{1*}^n$. Assuming that $\delta N \ll \zeta_1$ for the starting-point perturbations³, we find

$$\rho_{1,\text{ini}} \simeq \bar{\rho}_{1,\text{ini}} e^{3(1+w_{i,\text{ini}})\zeta_1}. \quad (29)$$

Although ϵ_w is approximately constant during evolution, one cannot ignore its perturbation. We find for $\epsilon_w \propto \phi_1^n$:

$$\epsilon_w = \epsilon_w^{(0)} + n\epsilon_w \frac{\delta\phi_1}{\bar{\phi}_1} + \dots \quad (30)$$

Here $n = 2$ for the quadratic potential in the slow-roll scenario.

In the calculation below we will omit the subscript “ini” when it is apparent. Substituting Eq.(28) into Eq.(29) we obtain for the inflating curvaton;

$$e^{3\epsilon_w \zeta_1} = \frac{V_0 \pm \frac{1}{2}m^2 \left[\bar{g} + \sum_{k=1}^{\infty} \frac{1}{k!} g^{(k)} \left(\frac{\bar{g}}{g'} \frac{\delta\phi_1}{\bar{\phi}_1} \right)^k \right]^2}{V_0 \pm \frac{1}{2}m^2 \bar{g}^2}, \quad (31)$$

where $(1 + w_{i,\text{ini}}) \equiv \epsilon_w$ is assumed for the inflating curvaton. Order by order, we have for the expansion $\rho_1 = \bar{\rho}_1 + \sum_{k=1} \frac{1}{k} \delta^{(k)} \rho_1$;

$$\delta^{(1)} \rho_1 = m^2 g \delta\phi_1 \quad (32)$$

$$\delta^{(2)} \rho_1 = m^2 \left(1 + \frac{gg''}{g'^2} \right) (\delta\phi_1)^2 \quad (33)$$

Defining the ratio $R \equiv \pm \frac{1}{2}m^2 \bar{\phi}_1^2 / \rho_1$, we find for $\zeta_1 \equiv \zeta_1^{(1)} + \frac{1}{2!}\zeta_1^{(2)} + \frac{1}{3!}\zeta_1^{(3)} + \dots$;

$$\begin{aligned} \zeta_1^{(1)} &\simeq \frac{2R}{3\epsilon_w} \frac{\delta\phi_1}{\bar{\phi}_1} \\ \zeta_1^{(2)} &= \frac{3\epsilon_w}{2R} \left(1 - 2R + \frac{gg''}{g'^2} - 4 \right) (\zeta_1^{(1)})^2. \end{aligned} \quad (34)$$

In the last line, -4 appears from the expansion of ϵ_w , which plays an essential role to get a signature of f_{NL} as will be seen later. In the slow-roll phase (A), we find

$$f_1 e^{3\epsilon_w (\zeta_1 - \delta N)} + (1 - f_1) e^{4(\zeta_2 - \delta N)} = 1. \quad (35)$$

Assuming $\zeta_2 \ll \delta N$ and expanding the equation, we obtain at first order

$$f_1 3\epsilon_w \left[\zeta_1^{(1)} - \delta N^{(1)} \right] + (1 - f_1) \left[-4\delta N^{(1)} \right] = 0, \quad (36)$$

where the expansion is given by $\delta N \equiv \delta N^{(1)} + \frac{1}{2!}\delta N^{(2)} + \frac{1}{3!}\delta N^{(3)} + \dots$. We can solve this equation as

$$\delta N^{(1)} = r_1 \zeta_1^{(1)} \simeq r_1 \zeta_{1,\text{ini}}^{\text{iso}(1)}. \quad (37)$$

At second order, we find

$$\begin{aligned} &-4(1 - f_1)\delta N^{(2)} + 16(1 - f_1)(\delta N^{(1)})^2 \\ &+ 3\epsilon_w f_1 \left(\zeta_1^{(2)} - \delta N^{(2)} \right) + 9\epsilon_w^2 f_1 \left(\zeta_1^{(1)} - \delta N^{(1)} \right)^2 \\ &+ 6\epsilon_w^{(1)} f(\zeta_1^{(1)} - \delta N^{(1)}) = 0, \end{aligned} \quad (38)$$

where the last line includes ϵ_w expansion. We thus find that

$$\begin{aligned} \delta N^{(2)} &= \left(\frac{3\epsilon_w f_1}{4(1 - f_1) + 3\epsilon_w f_1} \right) \left(\frac{1}{3\epsilon_w f_1} \right) (\delta N^{(1)})^2 \times \\ &\left[16(1 - f_1) + \frac{3\epsilon_w f_1}{r_1^2} \frac{\zeta_1^{(2)}}{(\zeta_1^{(1)})^2} + 9\epsilon_w^2 f_1 \left(\frac{1}{r} - 1 \right)^2 \right. \\ &\left. + \frac{6f_1}{r_1} \left(\frac{1}{r_1} - 1 \right) \frac{3\epsilon_w^2}{R} \right] \\ &= \left[\frac{1}{r_1} \frac{3\epsilon_w}{2R} \left(\frac{gg''}{g'^2} - 3 \right) + 3\epsilon_w r_1 \left(1 - \frac{1}{r_1} \right)^2 \right. \\ &\left. + \frac{16}{3\epsilon_w} \left(\frac{1-f}{f} \right) + \left(\frac{1}{r_1} - 1 \right) \frac{3\epsilon_w}{R} \right] (\delta N^{(1)})^2 \\ &= \left[\frac{1}{r_1} \frac{3\epsilon_w}{2R} \left(\frac{gg''}{g'^2} - 3 \right) - 6\epsilon_w + 4 + (3\epsilon_w - 4)r_1 \right. \\ &\left. \frac{3\epsilon_w}{r_1 R} + 3\epsilon_w \left(\frac{1}{r_1} - \frac{1}{R} \right) \right] \\ &\times (\delta N^{(1)})^2, \end{aligned} \quad (39)$$

where

$$r_1 = \frac{3\epsilon_w f_1}{4(1 - f_1) + 3\epsilon_w f_1} \quad (40)$$

has been used for the last equality. Our result reproduces the calculation of the standard curvaton when $\epsilon_w = 1$ (no expansion) and $R = 1$.

We thus find the non-Gaussianity parameter for the slow-rolling curvaton;⁴

$$f_{NL} = \frac{1}{r_1} \frac{5\epsilon_w}{4R} \left(\frac{gg''}{g'^2} - 1 \right)$$

³ $\delta N < \zeta_1$ is always needed for the curvaton mechanism. Otherwise the curvaton perturbation cannot dominate the spectrum.

⁴ This is partly different from earlier naive calculations [13].

$$-5\epsilon_w + \frac{10}{3} + \left(\frac{5}{2}\epsilon_w - \frac{10}{3}\right)r_1 + \frac{5}{2}\epsilon_w \left(\frac{1}{r_1} - \frac{1}{R}\right). \quad (41)$$

In contrast to the standard curvaton, a minus sign $R < 0$ is possible for the hilltop potential [23], which gives a positive f_{NL} . For the quadratic potential we find $\epsilon_w/R \sim \frac{2}{3}\eta_1 \equiv \text{const.}$ for the slow-roll curvaton, where $\eta_1 \equiv m^2/3H^2$ is the conventional slow-roll parameter during the curvaton inflation.⁵

One may choose either $\eta_1 \sim 1$ or $\eta_1 \ll 1$ for the specific model.

It is notable that f_{NL} can be expected to become the order of $O(10)$ in the inflating curvaton model. Although $\eta_1 \ll 1$ is expected in the radiation domination epoch [24], η_1 can become the order of unity during the curvaton inflation if the mass is not protected by the symmetry. When $\eta_1 \sim 1$ induces fast-roll inflation [25], the number of e-foldings is not large. e.g., $N_e \sim O(1)$ is expected in that case. Therefore we can expect $\frac{\epsilon_w}{R} \sim \eta_1 \sim 1$ and $r_1 \sim 0.1$ at the same time, which gives $f_{NL} \sim O(10)$ as a natural expectation of the current model,⁶ although the precise value of the non-Gaussianity depends on the initial conditions (more specifically on the “starting point”) and the details of the transition. In that sense, numerical calculation is mandatory for further study.

In the above calculation we have considered a simple quadratic potential. However, the curvaton potential during inflation may be given by a polynomial;

$$\rho_1 = V_0 + \frac{\lambda}{n} \frac{\phi_1^n}{M^{n-4}}, \quad (42)$$

and ϵ_w could have non-trivial ϕ_1 -dependence. In that case, assuming the instant decay after the slow-roll phase (A), we find for the inflating curvaton

$$\bar{\rho}_1 = V_0 + \frac{\lambda}{n} \frac{\phi_1^n}{M^{n-4}} \quad (43)$$

⁵ For the fast-roll scenario, ϵ_w will have a different η -dependence [32]. For the fast-roll field one will find $\tilde{c}H\dot{\phi}_1 \simeq V'$ with the coefficient defined by $\tilde{c} = (3 + \sqrt{9 - 12\eta_1})/2$, which gives $-H\frac{\delta\phi_1}{\phi_1} \simeq \frac{\tilde{c}}{3\eta_1} \frac{\delta\phi_1}{\phi_1}$. Our definition of ϵ_w ($\dot{\rho} = -3H\epsilon_w\rho$) can be combined with $\dot{\rho} \simeq V'\dot{\phi}_1$ to give $-H\frac{\delta\phi_1}{\phi_1} \simeq \frac{2}{3} \frac{R}{\epsilon_w} \frac{\delta\phi_1}{\phi_1}$. One may choose either $\tilde{c}$ or ϵ_w . The choice of the definition cannot cause any discrepancy in the result. The expansion of ϵ_w may not be trivial in the fast-roll scenario. See later section for the generalization.

⁶ The spectral index requires $\eta_1 \ll 1$ during primordial inflation, which must be explained in the specific inflationary model that creates the primordial perturbation $\delta\phi_1$ at the horizon exit. (Notice that we are using “ η_1 ” both in the primordial inflation stage and in the curvaton stage. The argument of the time is omitted.) Here we are expecting that the mechanism that suppresses η_1 during the primordial inflation is not valid during the curvaton inflation.

$$\rho_1 = V_0 + \frac{\lambda}{nM^{n-4}} \left[\bar{g} + \sum_{k=1} \frac{1}{k!} g^{(k)} \left(\frac{\bar{g}}{g'} \frac{\delta\phi_1}{\phi_1} \right)^k \right]^n. \quad (44)$$

Defining the ratio

$$R \equiv \frac{\frac{\lambda}{n} \frac{\bar{g}^n}{M^{n-4}}}{\bar{\rho}_1}, \quad (45)$$

and assuming $\epsilon_w \propto \phi^l$, we find order by order;

$$\zeta_1^{(1)} = \frac{nR}{3\epsilon_w} \frac{\delta\phi_1}{\phi_1} \simeq -H \frac{\delta\phi_1}{\dot{\phi}_1} \quad (46)$$

$$\frac{\zeta_1^{(2)}}{[\zeta_1^{(1)}]^2} = \frac{3\epsilon_w}{2R} \left[\frac{n(n-1)}{2} - \frac{n^2}{2}R + \frac{n}{2} \frac{gg''}{g'^2} - 2l \right] \quad (47)$$

In the slow-roll phase (A), the non-linear formalism gives Eq.(35). Expanding the equation, we obtain Eq.(36). At second order, we find

$$\begin{aligned} \frac{\delta N^{(2)}}{(\delta N^{(1)})^2} &= \left(\frac{3\epsilon_w f_1}{4(1-f_1) + 3\epsilon_w f_1} \right) \left(\frac{1}{3\epsilon_w f_1} \right) \\ &\times \left[16(1-f_1) + \frac{3\epsilon_w f_1}{r_1^2} \frac{\zeta_1^{(2)}}{(\zeta_1^{(1)})^2} \right. \\ &\quad \left. + 9\epsilon_w^2 f_1 \left(\frac{1}{r_1} - 1 \right)^2 + \frac{3f}{r_1} \left(\frac{1}{r_1} - 1 \right) \frac{3\epsilon_w^2 l}{2R} \right] \\ &= \frac{1}{r_1} \frac{3\epsilon_w}{2R} \left\{ \frac{n(n-1)}{2} + \frac{4-n^2}{2}R + \frac{n}{2} \frac{gg''}{g'^2} - 2l \right\} \\ &\quad + 3\epsilon_w r_1 \left(\frac{1}{r_1} - 1 \right)^2 + \frac{16}{3\epsilon_w} \left(\frac{1-f}{f} \right) \\ &\quad + \left(\frac{1}{r_1} - 1 \right) \frac{3\epsilon_w l}{2R} \\ &= \frac{1}{r_1} \frac{3\epsilon_w}{2R} \left\{ \frac{n(n-1)}{2} + \frac{4-n^2}{2}R + \frac{n}{2} \frac{gg''}{g'^2} - l \right\} \\ &\quad - 6\epsilon_w + 4 + (3\epsilon_w - 4)r_1 \\ &\quad + 3\epsilon_w \left(\frac{1}{r_1} - \frac{l}{2R} \right). \end{aligned} \quad (48)$$

Here the last equation is obtained using

$$f_1 = \frac{4r_1}{4r_1 + 3\epsilon_w(1-r_1)}. \quad (49)$$

We thus find the non-Gaussianity parameter

$$\begin{aligned} f_{NL} &= \frac{5\epsilon_w}{4r_1 R} \left[\frac{n(n-1)}{2} + \frac{4-n^2}{2}R + \frac{n}{2} \frac{gg''}{g'^2} - l \right] \\ &\quad + \frac{-15\epsilon_w + 10}{3} + \left(\frac{15\epsilon_w - 20}{6} \right) r_1 \\ &\quad + \frac{5\epsilon_w}{2} \left(\frac{1}{r_1} - \frac{l}{2R} \right). \end{aligned} \quad (50)$$

The standard curvaton corresponds to $n = 2$, $l = 0$, $R = 1$ and $\epsilon_w = 1$.

C. The Oscillating Curvaton

During the oscillating phase (B) we always find $w_1 = 0$. Therefore ζ_1 evaluated at the beginning of the oscillating phase (B) remain constant thereafter.

Considering the same calculation as in Eq.(11), we find

$$\delta N \equiv r_1 \zeta_1 + (1 - r_1) \zeta_2, \quad (51)$$

where in phase (B) we have

$$r_1 = \frac{3\bar{\rho}_1}{3\bar{\rho}_1 + 4\bar{\rho}_2}. \quad (52)$$

The “starting-point” quantities should be evaluated at the beginning of the oscillating phase (B). We thus find

$$\delta N_{\text{ini}} = \zeta_{\text{ini}} + [r_{1,\text{ini}} \zeta_{1,\text{ini}}^{\text{iso}} + (1 - r_{1,\text{ini}}) \zeta_{2,\text{ini}}^{\text{iso}}]. \quad (53)$$

Apart from the starting-point, the evolution of the curvature perturbation is caused by the time-dependent parameter r_1 and is given by

$$\delta N = \zeta_{\text{ini}} + r_1 \zeta_{1,\text{ini}}^{\text{iso}} + (1 - r_1) \zeta_{2,\text{ini}}^{\text{iso}}, \quad (54)$$

where the evolution of the curvature perturbation ($\delta N \neq \zeta_{\text{ini}}$) is possible due to the evolution of r_1 .

In the standard curvaton model, the beginning of the oscillating phase (B) is defined by the uniform density hypersurface $H \simeq m$, and the end is defined by the uniform density hypersurface $H \simeq \Gamma$. Therefore the modulation does not appear in the standard curvaton model. The deviation from the quadratic potential causes modulation at the beginning of the oscillation [26]. Note however that the deviation from the quadratic potential does not allow instant approximation at the transition, since the oscillation caused by the non-quadratic potential may not scale like matter immediately after the slow-roll phase. In that case the numerical study is essential [26].

Later in this paper we will calculate $\zeta_{1,\text{ini}}$ when there is modulation at the beginning of the oscillating phase. The calculation is essential when the hybrid curvaton [27] does not cause secondary inflation. In that case the hypersurfaces defining the end of the slow-rolling phase (uniform ρ_1 hypersurface defined by the waterfall) is significantly different from the uniform density hypersurfaces of the total density.

III. MODULATION IN THE CURVATON MECHANISM

A. Modulation in the non-linear formalism

The modulation at the transition ($w_{i-} \rightarrow w_{i+}$) can be implemented in the curvaton mechanism. Assuming that the transition occurs at the density ρ_x , the non-linear formalism can be separated as

$$\zeta_i = \delta N + \int_{\bar{\rho}_i}^{\rho_x} H \frac{d\tilde{\rho}_i}{3(1 + w_{i+})\bar{\rho}_i} + \int_{\rho_x}^{\rho_i} H \frac{d\tilde{\rho}_i}{3(1 + w_{i-})\bar{\rho}_i}$$

$$= \delta N + \frac{1}{3(1 + w_{i+})} \ln \left(\frac{\rho_x}{\bar{\rho}_i} \right) + \frac{1}{3(1 + w_{i-})} \ln \left(\frac{\rho_i}{\rho_x} \right) \quad (55)$$

Here $\bar{\rho}_i$ ($\rho_i > \rho_x > \bar{\rho}_i$) is placed after the transition so that it gives a well-defined starting-point for the curvaton mechanism after the transition.

In the pure curvaton mechanism we usually take the limit $\rho_x \rightarrow \rho_i$. (The transition is identical to the uniform density hypersurfaces.) In that case we can find ζ_i (for the transition from the slow-rolling phase (A) to the oscillating phase (B))

$$\zeta_i = \delta N + \frac{1}{3} \ln \left(\frac{\rho_i}{\bar{\rho}_i} \right), \quad (56)$$

which is identical to the conventional calculation of the curvaton mechanism.

On the other hand, for the hybrid curvaton [27] we must take the opposite limit $\rho_x \rightarrow \bar{\rho}_i$, where the critical point of the hybrid-type potential gives $\rho_x = \bar{\rho}_i = V(\phi_{i,c})$. In that case we can find (in contrast to the conventional curvaton);

$$\zeta_i = \delta N + \frac{1}{3\epsilon_w} \ln \left(\frac{\rho_i}{\bar{\rho}_i} \right), \quad (57)$$

which is enhanced when $\epsilon_w \ll 1$. Note that the enhancement is not due to the “evolution” (i.e, the curvaton mechanism), but caused by the “modulation” at the transition. In this case the evolution thereafter is nothing but the usual curvaton while the initial perturbation is enhanced by the “modulation”. Unlike the usual modulation, the above “modulation” is not due to the perturbation of an additional light field.

In contrast with the inflating curvaton, the hybrid curvaton [27] does not always assume the curvaton inflation. Then the “modulation” discussed above can play the crucial role.

This gives the new scenario of the modulation caused by the isocurvature perturbations. The source of the modulation is the isocurvature perturbation, which is identical to the conventional source of the curvaton mechanism. The gap between the uniform ρ_1 hypersurfaces and the uniform density hypersurfaces is the source of the modulation. Note however that in the single-component Universe there cannot be such modulation.

In contrast to the “modulation” discussed above, the “usual modulation” uses independent perturbation $\delta\rho_x \neq 0$ and the Universe is dominated by the single component. For instance, we know that

1. $\delta\Gamma \neq 0$ in the single-component Universe causes the modulated decay [15]
2. $\delta\phi_c \neq 0$ in the hybrid inflation model leads to the modulation of the critical point, but the density of the Universe is basically dominated by the single component. This scenario is sometimes called “the end of inflation” model [22].

In the next section we will discuss the usual modulation implemented in the multi-component Universe.

IV. MODULATION CAUSED BY THE INDEPENDENT MODULI

In this section we just consider the oscillating curvaton and the background radiation. The modulated decay of the curvaton is caused by the extra light field.

1. Modulated curvaton decay

If $\rho_2 > \rho_1$ before the decay, the radiation dominated Universe leads to $H^2 \propto a^{-4}$, which gives

$$\frac{\delta a_x}{a_x} = -\frac{1}{2} \frac{\delta \Gamma}{\Gamma}, \quad (58)$$

where a_x is the scale factor at the curvaton decay. The density perturbations created after the transition is

$$\begin{aligned} \frac{\delta \rho_1}{\rho_1} &\simeq 3(w_{1+} - w_{1-}) \frac{\delta a_x}{a_x} \\ &= -\frac{1}{2} \frac{\delta \Gamma}{\Gamma}. \end{aligned} \quad (59)$$

Therefore, the curvature perturbation generated by the modulation is

$$\begin{aligned} \zeta &\simeq \frac{1}{4} \frac{\delta \rho_1}{\rho} \\ &\simeq -\frac{1}{8} \frac{\rho_1}{\rho} \frac{\delta \Gamma}{\Gamma}. \end{aligned} \quad (60)$$

Here the factor $r \equiv \rho_1/\rho \ll 1$ can be used to enhance the non-Gaussianity, however the situation is completely the same as the conventional curvaton model.

2. Modulated decay (original scenario)

If $\rho_1 > \rho_2$ before the decay (matter dominated), we have $H^2 \propto a^{-3}$, which leads to

$$\frac{\delta a_x}{a_x} = -\frac{2}{3} \frac{\delta \Gamma}{\Gamma}. \quad (61)$$

This corresponds to the usual scenario of the modulated reheating (single component modulation), and gives

$$\zeta \simeq -\frac{1}{6} \frac{\delta \Gamma}{\Gamma}. \quad (62)$$

A. Modulation and curvaton after multi-field inflation

If the modulated decay and the multi-field inflation are the generic properties of the string theory, the modulation must be considered for the N-flation.

To illustrate our modulation mechanism in the typical multi-field inflation scenario, we consider a potential in which two inflaton fields have *the identical mass* [10]. The two-field chaotic inflation model has the symmetric potential given by

$$V(\phi_1, \phi_2) = \frac{1}{2} m^2 (|\phi_1|^2 + |\phi_2|^2) \equiv \frac{1}{2} m^2 |\phi_r|^2, \quad (63)$$

where we assumed that $\phi_{1,2}$ are complex scalar fields. We can define the angle θ defined as $\tan \theta = \frac{\phi_1}{\phi_2}$, which leads to the densities just at the end of inflation

$$\begin{aligned} \rho_{1,\text{end}} &= \frac{1}{2} m^2 M_p^2 \sin^2 \theta \\ \rho_{2,\text{end}} &= \frac{1}{2} m^2 M_p^2 \cos^2 \theta. \end{aligned} \quad (64)$$

In this model, the primordial perturbations are

$$\begin{aligned} \zeta^{\text{inf}} &\simeq \frac{1}{\eta} \frac{\delta \phi_{r*}}{\phi_{r*}} \simeq \frac{\sqrt{N_e}}{2\sqrt{6}\pi} \frac{m}{M_p} \\ \zeta_{1,\text{end}} &= \zeta^{\text{inf}} + \frac{2 \cos \theta}{3 \sin \theta} \delta \theta \\ \zeta_{2,\text{end}} &= \zeta^{\text{inf}} - \frac{2 \sin \theta}{3 \cos \theta} \delta \theta, \end{aligned} \quad (65)$$

where the fraction perturbation is sourced by the entropy perturbation δs during inflation. We find

$$\delta \theta = \frac{\delta s_*}{\phi_{r*}} = \frac{H_I}{2\pi \phi_{r*}} \simeq \frac{1}{4\pi \sqrt{6} N_e} \frac{m}{M_p}. \quad (66)$$

Besides the potential, which is nearly symmetric, we need the decay rate $\Gamma_1 \equiv \bar{\Gamma}_1 + \delta \Gamma_1$ and $\Gamma_2 \equiv \bar{\Gamma}_2 + \delta \Gamma_2$ for the fields.⁷ For simplicity we assume $\Gamma_2 > \Gamma_1$ and $\delta \Gamma_2 \ll \delta \Gamma_1$. In this section we assume $\rho_2 > \rho_1$ before the decay.

The component ϕ_2 decays at $H = \Gamma_2$, then ϕ_1 decays at $H = \Gamma_1$. When ϕ_1 decays the density ratio $f_1 \equiv \rho_1/\rho_{\text{total}}$ is given by

$$f_1|_{H=\Gamma_1} = \sin^2 \theta \sqrt{\frac{\Gamma_2}{\Gamma_1}}. \quad (68)$$

The creation of the curvature perturbation caused by the evolution (the curvaton mechanism) gives [10]

$$\begin{aligned} \delta N|_{\text{curv}} &\simeq f_1|_{H=\Gamma_1} \zeta_{1,\text{end}} \\ &\simeq \frac{2}{3} \sin \theta \cos \theta \sqrt{\frac{\Gamma_2}{\Gamma_1}} \delta \theta, \end{aligned} \quad (69)$$

⁷ The basic assumption behind this modulation is

$$\begin{aligned} \Gamma_i &= g_i(\varphi_j) m_i \\ &\simeq \bar{g}_i m_i + \left(\sum_j \frac{\partial g_i}{\partial \varphi_j} \delta \varphi_j \right) m_i + \dots, \end{aligned} \quad (67)$$

where φ_j is the moduli that determines the coupling constants in the effective action.

while the modulated decay creates the curvature perturbation;

$$\begin{aligned}\delta N|_{\text{mod}} &\simeq -\frac{1}{8}f_1|_{H=\Gamma_1}\frac{\delta\Gamma_1}{\Gamma_1} \\ &\simeq -\frac{\sin^2\theta}{8}\sqrt{\frac{\Gamma_2}{\Gamma_1}}\frac{\delta\Gamma_1}{\Gamma_1}.\end{aligned}\quad (70)$$

The domination by the modulated perturbation is possible if $\delta N|_{\text{mod}}$ dominates the curvature perturbation. From the above calculations we find that $\zeta^{\text{inf}} < \delta N|_{\text{mod}}$ gives

$$\frac{4\sqrt{N_e}}{\sqrt{6}\pi}\frac{1}{\sin^2\theta}\frac{m}{M_p}\sqrt{\frac{\Gamma_1}{\Gamma_2}} < \frac{\delta\Gamma_1}{\Gamma_1}, \quad (71)$$

while $\delta N|_{\text{curv}} < \delta N|_{\text{mod}}$ gives

$$\frac{4}{3\sqrt{6N_e}\pi}\frac{\cos\theta}{\sin\theta}\frac{m}{M_p} < \frac{\delta\Gamma_1}{\Gamma_1}. \quad (72)$$

Comparing the two conditions, the latter condition becomes significant when

$$N_e < \frac{1}{3}\sin\theta\cos\theta\sqrt{\frac{\Gamma_2}{\Gamma_1}}. \quad (73)$$

The non-Gaussianity parameter gives another bound for the modulation. Since the non-Gaussianity parameter f_{NL} is estimated as $f_{NL} \simeq \frac{3}{4f_1}$ when it is large, $f_{NL} \ll 10^2$ gives

$$\sin^2\theta\sqrt{\frac{\Gamma_2}{\Gamma_1}} \ll 10^2. \quad (74)$$

V. CONCLUSIONS AND DISCUSSIONS

In this paper we calculated the curvature perturbation for the generalized curvaton mechanism, including the curvaton mechanism caused by the slow-rolling curvaton. Our result shows that $f_{NL} \sim O(10)$ could be natural for the generalized curvaton mechanism.

In this scenario, modulation can be implemented in the curvaton mechanism. To explain the mechanism, the typical two-component Universe is considered, in which one component (curvaton) is slow-rolling. We first considered the new modulation scenario in which the source of the modulation is identical to the usual isocurvature perturbation that has been used for the conventional curvaton mechanism.

Then we considered the conventional modulation in the multi-component Universe⁸.

In this paper the modulation is specifically defined as the “inhomogeneous transition” in the curvaton sector, where the transition in the scaling of the density does not coincide with the uniform density hypersurfaces [14].

We examined the curvaton evolution and the modulation in the generalized curvaton scenarios, such as the inflating curvaton and the hybrid curvaton. For the slow-rolling curvaton, we calculated the non-Gaussianity using the non-linear formalism. This is the first calculation of the non-Gaussianity created during the slow-rolling phase of the curvaton. The relation between the curvaton evolution and the modulation has been revealed using the non-linear formalism.

For our purposes, we took the simplest set-ups for the calculation, which requires further study;

1. The instant phase transition is considered for simplicity, but the transition should be more complicated depending on the details of the model parameters [29]. In the standard curvaton scenario the sudden-decay approximation gives a good intuitive derivation, however in practice the curvaton density is continually decaying into radiation.
2. In this paper, we were avoiding the interaction between components [30]. If the interaction is significant the dissipation may appear, which (in the most extreme case) may lead to warm curvaton scenario [31].
3. In this paper the second component is always the radiation so that we can capture the essential of the scenario, however the assumption may not be valid in practice. We need to discuss multi-component Universe that could be a mixture of the slow-rolling, matter and radiation.
4. We have pointed out that the curvaton evolution and the “modulation” might be significant after conventional N-flation, however the model considered in this paper is based on the two-component Universe. We thus need some statistical argument for the evolution of the perturbations after N-flation.

Our calculation reveals the essential of the modulation and the generalized curvaton scenarios, giving intuitive arguments and the calculational details.

Note added: While finalizing this paper, we found a paper by M. Kawasaki, T. Kobayashi and F. Takahashi [32] which has some overlaps with our models.

VI. ACKNOWLEDGMENT

TM thanks D. H. Lyth and J. McDonald for many valuable discussions. This work is supported in part by Grant-in-Aid for Scientific research from the Ministry of Education, Science, Sports, and Culture (MEXT), Japan, No. 21111006, No. 22244030, and No. 23540327

⁸ When we are finishing this manuscript, a paper appears [28] but the modulation is “not” considered for the curvaton but for the inflaton.

(K.K.). S.E. is supported by the Grant-in-Aid for Nagoya University Global COE Program, "Quest for Fundamen-

tal Principles in the Universe: from Particles to the Solar System and the Cosmos".

-
- [1] D. H. Lyth, A. R. Liddle, "The primordial density perturbation: Cosmology, inflation and the origin of structure," Cambridge, UK: Cambridge Univ. Pr. (2009) 497 p.
 - [2] D. Wands, K. A. Malik, D. H. Lyth and A. R. Liddle, "A New approach to the evolution of cosmological perturbations on large scales," *Phys. Rev. D* **62**, 043527 (2000) [astro-ph/0003278].
 - [3] A. Gangui, F. Lucchin, S. Matarrese and S. Mollerach, "The Three point correlation function of the cosmic microwave background in inflationary models," *Astrophys. J.* **430**, 447 (1994) [astro-ph/9312033].
 - [4] M. Sasaki and E. D. Stewart, "A General analytic formula for the spectral index of the density perturbations produced during inflation," *Prog. Theor. Phys.* **95**, 71 (1996) [astro-ph/9507001].
 - [5] J. Garcia-Bellido and D. Wands, "Metric perturbations in two field inflation," *Phys. Rev. D* **53**, 5437 (1996) [astro-ph/9511029].
 - [6] V. F. Mukhanov and P. J. Steinhardt, "Density perturbations in multifield inflationary models," *Phys. Lett. B* **422**, 52 (1998) [astro-ph/9710038].
 - [7] D. Polarski and A. A. Starobinsky, "Isocurvature perturbations in multiple inflationary models," *Phys. Rev. D* **50**, 6123 (1994) [astro-ph/9404061].
 - [8] T. Matsuda, "Delta-N formalism for the evolution of the curvature perturbations in generalized multi-field inflation," *Phys. Lett. B* **682**, 163 (2009) [arXiv:0906.2525 [hep-th]]; T. Matsuda, "Elliptic Inflation: Generating the curvature perturbation without slow-roll," *JCAP* **0609**, 003 (2006) [hep-ph/0606137].
 - [9] F. Bernardeau and J. -P. Uzan, "NonGaussianity in multifield inflation," *Phys. Rev. D* **66**, 103506 (2002) [hep-ph/0207295]; L. E. Allen, S. Gupta and D. Wands, "Non-gaussian perturbations from multi-field inflation," *JCAP* **0601**, 006 (2006) [astro-ph/0509719]; H. Assadullahi, H. Firouzjahi, M. H. Namjoo and D. Wands, "Curvaton and the inhomogeneous end of inflation," arXiv:1207.7006 [astro-ph.CO].
 - [10] S. Enomoto and T. Matsuda, To appear
 - [11] T. Moroi, T. Takahashi, "Effects of cosmological moduli fields on cosmic microwave background," *Phys. Lett. B* **522**, 215-221 (2001). [hep-ph/0110096]. D. H. Lyth, D. Wands, "Generating the curvature perturbation without an inflaton," *Phys. Lett. B* **524**, 5-14 (2002). [hep-ph/0110002].
 - [12] Z. G. Berezhiani, A. S. Sakharov and M. Y. Khlopov, "Primordial background of cosmological axions," *Sov. J. Nucl. Phys.* **55**, 1063 (1992) [*Yad. Fiz.* **55**, 1918 (1992)]; A. S. Sakharov, D. D. Sokoloff and M. Y. Khlopov, "Large scale modulation of the distribution of coherent oscillations of a primordial axion field in the universe," *Phys. Atom. Nucl.* **59**, 1005 (1996) [*Yad. Fiz.* **59N6**, 1050 (1996)].
 - [13] K. Dimopoulos, K. Kohri, D. H. Lyth and T. Matsuda, "The inflating curvaton," *JCAP* **1203**, 022 (2012) [arXiv:1110.2951 [astro-ph.CO]].
 - [14] T. Matsuda, "Cosmological perturbations from an inhomogeneous phase transition," *Class. Quant. Grav.* **26**, 145011 (2009). [arXiv:0902.4283 [hep-ph]]; M. Kawasaki, T. Takahashi and S. Yokoyama, "Density Fluctuations in Thermal Inflation and Non-Gaussianity," *JCAP* **0912**, 012 (2009) [arXiv:0910.3053 [hep-th]].
 - [15] G. Dvali, A. Gruzinov and M. Zaldarriaga, "A new mechanism for generating density perturbations from inflation," *Phys. Rev. D* **69**, 023505 (2004) [arXiv:astro-ph/0303591].
 - [16] A. R. Liddle, A. Mazumdar, F. E. Schunck, "Assisted inflation," *Phys. Rev. D* **58**, 061301 (1998). [astro-ph/9804177]; S. Dimopoulos, S. Kachru, J. McGreevy, J. G. Wacker, "N-flation," *JCAP* **0808**, 003 (2008). [hep-th/0507205].
 - [17] A. D. Linde and V. F. Mukhanov, "Nongaussian isocurvature perturbations from inflation," *Phys. Rev. D* **56** (1997) 535 [arXiv:astro-ph/9610219].
 - [18] D. H. Lyth, "Can the curvaton paradigm accommodate a low inflation scale?," *Phys. Lett. B* **579** (2004) 239 [hep-th/0308110].
 - [19] M. Sasaki, J. Valiviita and D. Wands, "Non-Gaussianity of the primordial perturbation in the curvaton model," *Phys. Rev. D* **74**, 103003 (2006) [astro-ph/0607627].
 - [20] D. H. Lyth, K. A. Malik and M. Sasaki, "A General proof of the conservation of the curvature perturbation," *JCAP* **0505**, 004 (2005) [astro-ph/0411220];
 - [21] D. Langlois, F. Vernizzi and D. Wands, "Non-linear isocurvature perturbations and non-Gaussianities," *JCAP* **0812**, 004 (2008) [arXiv:0809.4646 [astro-ph]].
 - [22] D. H. Lyth, "Generating the curvature perturbation at the end of inflation," *JCAP* **0511**, 006 (2005) [astro-ph/0510443].
 - [23] T. Matsuda, "Hilltop curvatons," *Phys. Lett. B* **659**, 783 (2008) [arXiv:0712.2103 [hep-ph]].
 - [24] D. H. Lyth and T. Moroi, "The Masses of weakly coupled scalar fields in the early universe," *JHEP* **0405**, 004 (2004) [hep-ph/0402174].
 - [25] A. D. Linde, "Fast roll inflation," *JHEP* **0111**, 052 (2001) [hep-th/0110195].
 - [26] M. Kawasaki, T. Kobayashi and F. Takahashi, "Non-Gaussianity from Curvatons Revisited," *Phys. Rev. D* **84** (2011) 123506 [arXiv:1107.6011 [astro-ph.CO]].
 - [27] K. Dimopoulos, K. Kohri and T. Matsuda, "The hybrid curvaton," *Phys. Rev. D* **85**, 123541 (2012) [arXiv:1201.6037 [hep-ph]].
 - [28] K. -Y. Choi and O. Seto, "Modulated reheating by curvaton," arXiv:1204.1419 [astro-ph.CO].
 - [29] T. Matsuda, "Cosmological perturbations from inhomogeneous preheating and multi-field trapping," *JHEP* **0707**, 035 (2007) [arXiv:0707.0543 [hep-th]].
 - [30] T. Matsuda, "Modulated Inflation," *Phys. Lett. B* **665**, 338 (2008) [arXiv:0801.2648 [hep-ph]]; T. Matsuda, "Modulated inflation from kinetic term," *JCAP* **0805**, 022 (2008) [arXiv:0804.3268 [hep-th]].
 - [31] T. Matsuda, "Entropy production and curvature perturbation from dissipative curvatons," *JCAP* **1009** (2010) 006 [arXiv:1007.3636 [astro-ph.CO]]; T. Matsuda, "As-

- pects of warm-flat directions,” *Int. J. Mod. Phys. A* **25**, 4221 (2010) [arXiv:0908.3059 [hep-ph]]; T. Matsuda, “Evolution of the curvature perturbations during warm inflation,” *JCAP* **0906**, 002 (2009) [arXiv:0905.0308 [astro-ph.CO]].
- [32] M. Kawasaki, T. Kobayashi and F. Takahashi, “Non-Gaussianity from Axionic Curvaton,” arXiv:1210.6595 [astro-ph.CO].